

International Publication Number: WO 00/09518

International Application Number: PCT/JP99/04256

Filing Date of International Application: August 5, 1999



|   |   |   |
|---|---|---|
| (51) 国際特許分類6<br>C07F 9/08, 9/6581, C08J 5/00, C08K<br>5/5399, C08L 85/02, 101/00, C09K<br>21/12, 21/14  | A1  | (11) 国際公開番号<br>WO00/09518<br><br>(43) 国際公開日<br>2000年2月24日(24.02.00) |
| (21) 国際出願番号<br>PCT/JP99/04256<br><br>(22) 国際出願日<br>1999年8月5日(05.08.99)<br><br>(30) 優先権データ<br>特願平10/228897 1998年8月13日(13.08.98)<br>特願平10/248415 1998年9月2日(02.09.98)  | (74) 代理人<br>弁理士 三枝英二, 外(SAEGUSA, Eiji et al.)<br>〒541-0045 大阪府大阪市中央区道修町1-7-1<br>北浜TNKビル Osaka, (JP)   |   |
| (71) 出願人 (米国を除くすべての指定国について)<br>大塚化学株式会社 (OTSUKA CHEMICAL CO., LTD.)[JP/JP]<br>〒540-0021 大阪府大阪市中央区大手通3丁目2番27号<br>Osaka, (JP)<br>(72) 発明者; および<br>(75) 発明者/出願人 (米国についてのみ)<br>中長偉文(NAKACHO, Yoshifumi)[JP/JP]<br>荻原忠男(YABUHARA, Tadao)[JP/JP]<br>多田祐二(TADA, Yuji)[JP/JP]<br>西岡洋一(NISHIOKA, Yoichi)[JP/JP]<br>〒771-0130 徳島県徳島市川内町加賀須野463<br>大塚化学株式会社 徳島研究所内 Tokushima, (JP) | (81) 指定国 AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU,<br>CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR,<br>LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR,<br>TT, UA, US, UZ, VN, YU, ZA, 欧州特許 (AT, BE, CH, CY, DE,<br>DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI特許<br>(BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD,<br>TG), ARIPO特許 (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW),<br>ユーラシア特許 (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM)<br><br>添付公開書類<br>国際調査報告書 |   |

(54) Title: CROSS-LINKED PHENOXYPHOSPHAZENE COMPOUND, PROCESS FOR PRODUCING THE SAME, FLAME  
 RETARDANT, FLAME-RETARDANT RESIN COMPOSITION, AND MOLDED FLAME-RETARDANT RESIN

(54) 発明の名称 架橋フェノキシホスファゼン化合物、その製造法、難燃剤、難燃性樹脂組成物及び難燃性樹脂成形体

(57) Abstract

A flame retardant which has advantages, for example, that it is halogen-free, has a high melting point, is lowly volatile, and does not impair the properties inherent in resins. The flame retardant comprises a cross-linked phenoxyphosphazene compound which is a compound obtained by cross-linking a phosphazene compound with cross-linking groups, e.g., phenylene groups, each cross-linking group being interposed between two oxygen atoms left by phenyl group elimination, and which has no free hydroxyl group (derived from the cross-linking group) in the molecule and contains phenyl groups in an amount of 50 to 99.9 % based on the total number of all the phenyl groups in the phosphazene compound.